

Book of abstracts of the 4th CHRC Annual Summit

Abstracts

Évora, Portugal, May 25 and 26, 2023

Editorial Board

Jorge Bravo: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal.
Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (jorgebravo@uevora.pt)

Rute Dinis de Sousa: NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC) Universidade NOVA de Lisboa (NMS/UNL), Lisboa, Portugal. (rute.sousa@nms.unl.pt)

Gabriela Almeida: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal.
Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (gsna@uevora.pt)

Dulce Cruz: Escola Superior de Enfermagem São João de Deus, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (dcruz@uevora.pt)

Otilia Zangão: Escola Superior de Enfermagem São João de Deus, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (otiliaz@uevora.pt)

Bruno Gonçalves: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (bgoncalves@uevora.pt)

Rui Raimundo: Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (rjcr@uevora.pt)

Nadiya Bocharova: Comprehensive Health Research Centre (CHRC), Universidade NOVA de Lisboa, Lisboa, Portugal. (nadiya.bocharova@nms.unl.pt)

Andreia Santos: Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (andreia.santos@uevora.pt)

Lúcia Domingues: NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC) Universidade NOVA de Lisboa (NMS/UNL), Lisboa, Portugal. (lucia.domingues@nms.unl.pt)

Margarida Santos Dias: Institute of Global Mental Health, Lisbon, Portugal. Comprehensive Health Research Centre (CHRC), Lisbon, Portugal. (margarida.dias@nms.unl.pt)

Ana Rita Pedro: National School of Public Health, Public Health Research Centre, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), National School of Public Health, Lisboa, Portugal. (rita.pedro@ensp.unl.pt)

Hugo Folgado: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (hfolgado@uevora.pt)

Armando Raimundo: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal. (ammr@uevora.pt)

Contact editor

Jorge Bravo (jorgebravo@uevora.pt)

Executive Committee Members

Ana Rodrigues

Armando Raimundo

Helena Canhão

Rute Dinis de Sousa

Sónia Dias

Scientific Committee Members (Reviewers)

Ana Filipa Mourão: NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), Universidade NOVA de Lisboa, Lisboa, Portugal.

Carla Martins: National School of Public Health, Public Health Research Centre, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), National School of Public Health, Lisboa, Portugal.

Carla Lança: National School of Public Health, Public Health Research Centre, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), National School of Public Health, Lisboa, Portugal.

Catarina Pereira: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal.

Diana Teixeira: NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), Universidade NOVA de Lisboa, Lisboa, Portugal.

João Pedro Santos: National School of Public Health, Public Health Research Centre, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), National School of Public Health, Lisboa, Portugal.

Joana Alves: National School of Public Health, Public Health Research Centre, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), National School of Public Health, Lisboa, Portugal.

Maria Luz Barros: Escola Superior de Enfermagem São João de Deus, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal.

Nuno Mendonça: NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), Universidade NOVA de Lisboa, Lisboa, Portugal.

Orlando Fernandes: Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal.

Pedro Martins: NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal. Comprehensive Health Research Centre (CHRC), Universidade NOVA de Lisboa, Lisboa, Portugal.

Susana Mendonça: Escola Superior de Enfermagem São João de Deus, Universidade de Évora, Évora, Portugal. Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal.

Disclosure Statement

The authors and editorial board elements declare no conflict of interest.

Funding

This work is funded by national funds through the Foundation for Science and Technology, under the project UIDB/04923/2020.

S. Karger
Medical and Scientific Publishers

Disclaimer

The statements, opinions and data contained in this publication are solely those of the individual authors and contributors and not of the publisher and the editor(s). The appearance of advertisements in the journal is not a warranty, endorsement, or approval of the products or services advertised or of their effectiveness, quality or safety. The publisher and the editor(s) disclaim responsibility for any injury to persons or property resulting from any ideas, methods, instructions or products referred to in the content or advertisements.

Drug Dosage

The authors and the publisher have exerted every effort to ensure that drug selection and dosage set forth in this text are in accord with current recommendations and practice at the time of publication. However, in view of ongoing research, changes in government regulations, and the constant flow of information relating to drug therapy and drug reactions, the reader is urged to check the package insert for each drug for any change in indications and dosage and for added warnings and precautions. This is particularly important when the recommended agent is a new and/or infrequently employed drug.

karger@karger.com
www.karger.com/pjp

Karger 

All rights reserved.

No part of this publication may be translated into other languages, reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, recording, microcopying, or by any information storage and retrieval system, without permission in writing from the publisher or, in the case of photocopying, direct payment of a specified fee to the Copyright Clearance Center (see 'General Information').

© Copyright 2024 NOVA National School of Public Health
Published by S. Karger AG, Basel,
P.O. Box, CH-4009 Basel (Switzerland)

Evaluating Insulin and C-Peptide Levels: How Method and Brand Affect Results

Maria João Meneses^{1,2}, Rita S. Patarrão¹, Artur Romão²,
João Nabais³, João F. Raposo^{1,4}, M Paula Macedo^{1,4}

¹iNOVA4Health, NOVA Medical School|Faculdade de Ciências Médicas, NMS|FCM, Universidade NOVA de Lisboa; Lisboa, Portugal; ²DECSIS II Iberia, Évora, Portugal; ³Comprehensive Health Research Centre (CHRC), Departamento de Ciências Médicas e da Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal; ⁴APDP – Diabetes Portugal, Education and Research Center, Lisbon, Portugal

Keywords: C-Peptide; Diabetes; Evaluation methods; Insulin
Objective: Throughout the past several decades, diabetes diagnosis and treatment have focused on glycemia and glycated hemoglobin. However, it has become apparent that the evaluation of other biochemical indicators, such as insulin and C-peptide, provides a broader view of each individual's metabolic status, enabling a precision medicine approach since diabetes therapy goes beyond glucose control. Our aim was to assess the differences, if any, on C-peptide and insulin levels' results when using different methods and/or the same method from different vendors, to evaluate the reproducibility of results throughout the distinct approaches.

Methods: Participants recruited at the APDP clinic underwent a 75 g OGTT, and blood samples were collected at baseline, 30, 90, and 120 minutes after the glucose load. Insulin and C-peptide were measured by chemiluminescence and three commercially available ELISA kits from different brands (A, B, and C).

Results: Using the chemiluminescence assay for both insulin and C-peptide resulted in higher levels than ELISAs ($p<0.05$). Regarding insulin, there were discrepancies in the results obtained both between the different ELISAs and the chemiluminescence method. Although in the detection range, ELISA C did not perform well below 40 μ IU/mL. In contrast, ELISA A was very sensitive, and a dilution was required. In general, there were also discrepancies in C-peptide results, but less than with insulin. Overall, ELISA A seemed to be the most reliable. ELISA B is the only one that shows peaks at 90 min, with all the other three following the same trend between them, although with slightly different levels: Chemiluminescence presents the highest values, followed by ELISA A and then by ELISA C.

Conclusions: In conclusion, we found that different methods and vendors do not provide consistent results between them, and consequently, the evaluation of results may be biased. This should be considered when using these values for disease management evaluation or further used in indexes of insulin secretion, metabolism, and/or resistance.

Depression in older adults during the pandemic by COVID-19: Rethink the future

Celso Silva^{1,2,3}, César Fonseca^{2,3}, Rogério Ferreira^{1,3},
Lara Pinho^{2,3}

¹Higher School of Health, Polytechnic Institute of Beja, Beja, Portugal; ²Nursing Department, University of Évora, Évora, Portugal; ³Compreensível Health Research Center, Universidade de Évora, Évora, Portugal

Keywords: Crisis; Depressive symptoms; Health policy; Older adults, Pandemic consequences

Objective: To describe the determinants of depressive symptoms in adults aged 60 years and older related with the COVID-19 pandemic and contribute to the definition of health policies in similar situations in the future.

Methods: This study adopts a systematic literature review and the search was conducted in several databases to identify eligible studies that were published between December 2019 and March 2022. The inclusion criteria were as follows: Studies in which participants were diagnosed with or screened for a depressive disorder, regardless of progression status and the presence or absence of other conditions, in older adults aged 60 years or older. The review includes studies on factors associated with depression in older adults since the onset of the pandemic by COVID-19, regardless of geographical location and context (community, culture or specific environment). The review includes primary quantitative empirical studies: cross-sectional, longitudinal, observational or experimental studies. This review also includes studies with or without a comparison group. Search terms included combinations of the following three subject headings according to Medical Subject Headings (MeSH) terms: Depression, Depressive Symptoms, Older Adult or COVID-19. Articles with the following terms or combinations or terms in the title, keywords and/or abstract were retained: ("Depression") OR ("Depressive Disorder") OR ("depressive symptoms") AND ("Older Adults") OR ("Aging") OR ("Elderly") AND ("Pandemic") OR (COVID-19).

Results: Fifty-three studies were included, seventeen of which identified determinants of depression directly associated with the pandemic COVID-19, while the remainder identified determinants of depression already widely known. Regarding the determinants directly associated with the pandemic COVID-19 we grouped the results into 4 categories: 1) Factors and feelings related to stress or worries related to the COVID-19 pandemic: Distress related to COVID-19, the feeling of being very vulnerable to the risk of contracting COVID-19, the fear of infecting others and feelings related to the high mortality due to COVID-19. 2) Factors related to receiving news about COVID-19 in the media: Receiving news about COVID-19 through the media,

having insufficient knowledge about the pandemic and receiving COVID-19 related information from health workers. 3) Factors related to measures that were taken to reduce the spread of COVID-19: Containment measures, COVID-19 related social agitation, difficulty receiving healthcare, difficulty obtaining medication. 4) Factors related to concerns about family, friends, and acquaintances: Existence of infected acquaintances and the loss of family and friends.

Conclusions: A better understanding of the pandemic-associated factors that may determine the onset of depression in this population may be helpful in reassessing and rethinking health responses in a future pandemic situation. There needs to be a balance between the need to contain the virus and the concern for maintaining the mental health of older adults.

Rehabilitation technologies with the potential to be incorporated into the clinical practice of physiotherapists: a systematic review

Tarciano Siqueira^{1,2}, José Parraça^{1,2}, João Paulo Sousa^{1,2}

¹ Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre (CHRC), Universidade de Évora, Évora, Portugal

Keywords: Physiotherapy; Rehabilitation; Health technologies; Artificial intelligence; Technological rehabilitation

Objective: This review seeks to identify and synthesise the best available evidence on technologies used for the management of physical rehabilitation in patients, with the potential to be integrated into the physiotherapist's clinical practise.

Methods: We carried out a prospective record of the protocol for carrying out this systematic review that was published on the PROSPERO platform. The guidelines of the list of PRISMA recommendations were considered following the guidelines of the updated guidance and examples for reporting systematic reviews. A search was carried out through the databases Embase, PubMed, Physiotherapy Evidence Database (PEDro) and Ovid, carried out in the time period from 2000 to 2020, published until 2021 in Portuguese or English.

Results: There were a total of 121 publications found, 18 of which were fully selected randomized studies for analysis and inclusion. In the included studies, the use of technology during various phases of the clinical approach to rehabilitation was evident. The studies that met the inclusion criteria were classified according to the type of technology and its role in the clinical approach to rehabilitation, namely digital technologies; Artificial Intelligence and Robotics; virtual technologies or hybrid technologies, given that their composition includes more than one of the aforementioned types of technology.

Conclusions: Rehabilitation technologies can effectively mediate clinical activities of the physiotherapist, such as preventing injuries, monitoring movements, and coordinating rehabilitation programmes, with minimal or no intervention from the physiotherapist. More research is required to demonstrate with greater specificity the potential for different technologies to assist the physiotherapist in the curative or preventative care of patients' physical health.

Health policies and public health and social measures impact measures on COVID-19 in Europe: a review and modified Delphi technique

Marília Silva Paulo¹, Mariana Peyroteo¹, Melanie Maia², Luís Velez Lapão^{1,2,3}

¹Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal; ²UNIDEMI, NOVA School of Science and Technology, Universidade Nova de Lisboa, Portugal; ³LASI, Intelligent Systems Associated Laboratory, Guimarães, Portugal

Keywords: COVID-19; Europe; health policy; Public Health and Social Measures; Non-pharmaceutical interventions

Objective: This study aimed at reviewing the literature regarding the key results of the public health and social measures (PHSM) implemented during COVID-19 pandemic, and it identified the three PHSM considered to have most impacted the epidemiological curve of COVID-19 over the last two years, in different stages of the pandemic.

Methods: The PHSM under study were selected from the Oxford Stringency Index Government Response Tracker (OxCGRT) combined with the topics presented during the Rapid Exchange Forum (REF) meetings in the scope of the Population Health Information Research Infrastructure (PHIRI) project (H2020). A Rapid Review was conducted using Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines to identify which reviews have already been published about each PHSMs and their results. In addition, two Modified Delphi Panel surveys were conducted to uphold the results found.

Results: There were 3212 studies retrieved from PubMed and 162 full texts assessed for eligibility and 37 were included in this PHSMs summary. The three most significant and evidence-based measures from the review were hygiene measures, mask measures and vaccination policies. From the modified Delphi Panel, the most priority PHSMs identified on the four waves in questions were case isolation at home, face coverings, testing policy and social distancing, respectively.

Conclusions: The evidence found is very important for both researcher and policy-makers. The study of PHSM's impact on COVID-19 is significant as a lesson learned for the expected next pandemic and as a contribution to the health systems resilience discussion. These lessons should also be considered in both educational and preparedness programs as well as the understanding from the perspectives of the subject matter experts and the scientific evidence.

Patients' Voice in Breast Cancer Surgery Through PROM and PREM

Anabela Coelho^{1,2,3}, Candan Kendir⁴, Eliana Barreiro⁴ and Tânia Gaspar^{5,6}

¹Comprehensive Health Research Centre (CHRC), Nursing Department, University of Évora, 7004-516 Évora, Portugal;

²H&TRC-Health & Technology Research Center, ESTeSL-Escola Superior de Tecnologia da Saúde, Instituto Politécnico de Lisboa, 1549-020 Lisbon, Portugal; ³Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade NOVA de Lisboa, 1099-085 Lisbon, Portugal; ⁴Organisation for Economic Co-Operation and Development, 75016 Paris, France; ⁵Instituto de Saúde Ambiental (ISAMB), Faculdade de Medicina, Universidade de Lisboa, 1649-028 Lisbon, Portugal; ⁶Digital Human-Environment Interaction Labs (HEI-LAB), Universidade Lusófona de Humanidades e Tecnologias, 1749-024 Lisbon, Portugal

Keywords: Patient Outcomes; Quality in Health; PROM; PREM; Health System

Objective: The aim of this study is to measure breast surgery outcomes reported by Portuguese patients.

Methods: The current study was a prospective transversal study conducted by the research team from Évora University and Lusiadas University in Portugal. Data were anonymously collected by 10 public hospitals from National Health Service from March 2021 to March 2022. The data collection procedure and analysis followed the "OECD Breast Cancer Patient Reported Outcomes Working Group" protocol, allowing subsequent comparability with data from other OECD member countries. The quality of life, as a complementary variable, was measured through the reduced version of the 6-item WHOQoL-brief scale.

Results: The study included 378 women with breast cancer, with the age distribution being 19.8% aged 15 to 49 years and 80.2% aged 50 years and over. Most women were satisfied with the treatment outcome regarding the shape of their lumpectomy breast when wearing a bra (96.1%) and with the equal size of both breasts (78.3%). Obese women (p-value= 0.006) and women whom did radiotherapy (p-value=0.028) show less satisfaction with the treatment.

Conclusions: Measuring the outcomes and experiences is a fundamental step for the continuous improvement of quality in healthcare. Measuring breast cancer outcomes and experiences from Portuguese patients' perspectives gives us insight into the quality-of-care services and reinforces that patients' satisfaction and quality of life depended on multiple factors like the existing previously to the treatment (e.g. obesity); factors related to treatment (e.g. radiotherapy; breast asymmetry), and also factors that were unrelated with treatment (e.g. social relationships).

Fungal and Bacterial Infections Discrimination in ICU Patients based on serum molecular fingerprint

Rúben Araújo^{1,2}, Luís Ramalhete^{3,4}, Tiago Fonseca², Cristiana Von Rekowski², Luís Bento^{1,5,6}, Cecília R.C. Calado²

¹Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal; ²Laboratory of Health and Engineering, Instituto Superior de Engenharia de Lisboa, Portugal; ³Blood and Transplantation Center of Lisbon, Instituto Português do Sangue e da Transplantação, Alameda das Linhas de Torres, nº 117, 1769 – 001 Lisboa, Portugal; ⁴iNOVA4Health - Advancing Precision Medicine, RG¹¹: Reno-Vascular Diseases Group, NOVA Medical School, Faculdade de Ciências Médicas, Universidade NOVA de Lisboa, 1169-056 Lisbon, Portugal; ⁵Intensive Care Department, Centro Hospitalar Universitário de Lisboa Central, CHULC; Rua José António Serrano, 1150-199 Lisboa, Portugal; ⁶Integrated Pathophysiological Mechanisms, CHRC, NOVA Medical School, Faculdade de Ciências Médicas, NMS, FCM, Universidade NOVA de Lisboa; Campo Mártyres da Pátria, 130, 1169-056 Lisboa, Portugal

Keywords: FTIR spectroscopy; Intensive Care Unit; Fungal infection; Bacterial infection, Diagnostics

Objective: The rapid discrimination between fungal and bacterial infections in intensive care unit (ICU) patients is crucial for an efficient antimicrobial therapy. This work aims to develop a predictive model capable of discriminating fungal and bacterial infections among these patients based on a rapid serum analysis.

Methods: Serum from 46 patients at ICU, with COVID-19, were analyzed by Fourier-Transform infrared (FTIR) spectroscopy: 17 did not present any other infection; 12 presented bacterial infections (including *Enterobacter aerogenes* (2), *Escherichia coli* (4), *Enterococcus faecalis* (1), *Haemophilus influenzae* (1) and *Serratia marcescens* (1), *Klebsiella pneumoniae* (3)), isolated from blood, tracheobronchial aspiration, bronchoalveolar lavage and urine; 17 with fungal infections (including *Candida albicans* (14), *C. tropicalis* (1), *C. parapsilosis* (1) and *C. glabrata* (1), isolated from bronchoalveolar lavage, urine and urethral exudate. The impact of a Fast Correlation Based Filter (FCBF) of normalized second derivatized spectra (between 406-1800 and 2800-3992 cm⁻¹) was evaluated on a t-distributed Stochastic Neighbor Embedding (t-SNE), and a Naïve-Bayes model. The predictive models were based on 10 random iterations, with 80% of samples used for training and the remaining 20% for validation.

Results: The impact of FCBF was evaluated on t-SNE to select the spectral bands with high significance in detecting the infection and in discriminating the bacterial from the fungus infection. Optimized Naïve-Bayes models enabled to detect the infection with a sensitivity of 83% and a specificity of 81%, and the discrimination of the bacterial and fungal infection with a sensitivity and specificity of 86% and 96%, respectively.

Conclusions: Serum analysis, based on FTIR spectroscopy associated to machine learning algorithms presents a high potential to detect in a rapid and economic mode either the infection either the discrimination between bacterial or fungal infection.

The association of Internet use and health related quality of life in older adults: longitudinal analysis of a population-based cohort

Ana Rita Henriques¹, Nuno Mendonça¹, Helena Canhão¹,
Ana M Rodrigues¹

¹Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal

Keywords: Older adults; Internet use; Quality of life

Objective: The relationship between internet use and the health of older adults is still unclear. Therefore, we aimed to assess the impact of internet use on quality of life and physical function among Portuguese older adults (60+ years) using a prospective analytical approach.

Methods: This study included participants aged 60 or over from a nationwide population-based cohort (EpiDoC Cohort) and followed them up to 7 years. The second wave of the EpiDoC cohort was used as the baseline. Being an internet user, the years as user, and length of usage per/week was self-reported. Health-related quality of life (HRQoL) was measured using the EuroQol 5-dimensions questionnaire, physical function was assessed using the Health Assessment Questionnaire. Linear mixed model and random intercept tobit regression were used to assess the association between internet usage with HRQoL and physical function, respectively. A sensitivity analysis was performed considering years as an internet and frequency of use.

Results: This study followed 2272 participants, of which 20.9% reported using internet for an average of 9.5 years. The average internet usage per week was 8.2 days, and 4.9% of users reported using it for purposes related to their health. There was an association between internet users ($\beta=0.06$ [0.02; 0.09]; $\beta=-0.15$ [-0.25; -0.04]) and improved HRQoL and physical function over time, respectively, when compared with non-users, adjusting for time from baseline, sex, age, education level, BMI, smoking habits, alcohol consumption, regular exercise and multimorbidity.

Conclusions: Our results suggest that internet use among older adults is associated with an improvement in HRQoL and physical function. Thus, investing in digital literacy and digital solutions becomes essential for this population. Sensitivity analysis revealed that being an experienced internet user vs. a non-user ($\beta=0.07$ [0.03; 0.12]; $\beta=-0.24$ [-0.38; -0.10]) was associated with improved HRQoL and physical function over time, respectively.

Trajectories and determinants of Ageing in Portugal: Insights from EpiDoC, a nationwide population-based cohort

David G. Lopes¹, Nuno Mendonça¹, Ana Rita Henriques¹,
Jaime Branco^{1,2}, Helena Canhão^{1,2}, Ana M Rodrigues^{1,3}

¹EpiDoC Unit, CHRC, NOVA Medical School, Universidade NOVA de Lisboa, Lisboa, Portugal; ²Centro Hospitalar Lisboa Ocidental (CHLO-E.P.E.), Serviço de Reumatologia do Hospital Egas Moniz, Portugal; ³Rheumatology Unit, Hospital dos Lusíadas, Portugal

Keywords: Disability; Quality of life; Older adults; Trajectories; Longitudinal

Objective: Portugal's population is ageing due to increased life expectancy and reduced fertility rates. We aimed to estimate the health trajectories of the Portuguese older adults in a 10-year period and to assess associated sociodemographic, lifestyle factors and multimorbidity status.

Methods: Using data from the population-based EpiDoC cohort, we estimated, using linear mixed models, the trajectories of quality of life and physical function over 10 years in 4135 Portuguese older adults. Associated factors were assessed using linear mixed models (quality of life) and random intercept tobit regression (physical function).

Results: Physical disability of participants increased by 0.26 (0.24, 0.29), and quality of life declined by 0.07 (-0.08, -0.06), over 10 years. With advancing age, older adults reported a faster reduction in quality of life and faster increase in physical disability. At baseline, 24.8% of older adults reported a lot of difficulty or were unable to perform activities in dressing, arising, eating, and bathing. In general, women were in worse health than men at baseline, albeit with a similar rate of change throughout the follow-up. Higher education, alcohol consumption and regular exercise were associated with better quality of life and physical function while multimorbidity and excess weight were associated with worse reporting of these outcomes.

Conclusions: These findings, based on longitudinal data with 10 years of follow-up, are essential to effectively plan resource allocation, plan better healthcare and design informed public health policies in Portugal.

Family-Centered Care Effects in Family with Child(ren) with Intellectual Disability: A Review

Teresa Mestre^{1,2}, Ermelinda Caldeira¹, Manuel José Lopes¹

¹Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ²Escola Superior de Saúde, Instituto Politécnico de Beja, Portugal

Keywords: Family Nursing; Health Promotion; Intellectual Disability; Outcome Assessment, Health Care; Systematic Review

Objective: To determine what effects or health gains result from family-centered care (FCC) in families with child(ren) with intellectual disabilities and identify its components considered effective.

Methods: A systematic review of quantitative studies was conducted. MEDLINE with Full Text, CINAHL PLUS with Full Text,

Academic Search Complete and Psychology and Behavioral Sciences Collection databases were searched through EBSCOhost Web platform. The PRISMA guidelines were applied, and the protocol of the study was registered with PROSPERO (CRD42023398902). The quality of the included studies was appraised using the STrengthening the Reporting of OBservational Studies in Epidemiology (STROBE) guidelines for cross-sectional studies.

Results: Eight studies were selected and the study of the perception's and involvement of family in care was the main source of outcomes on FCC. It was identified as main components: collaboration between family members and healthcare professionals; assessment of the family context; and health policies/ access to healthcare services.

Conclusions: FCC is widely used in the paediatric context, but its implementation remains a challenge for health services. The effects of FCC refer to the promotion of family well-being and self-care; family satisfaction and consequent evaluation of the care process; and family readaptation. The needs and expectations arising from the family perception should be studied to provide a more effective response to the FCC.

AlenRiscos Observatory: tobacco, alcohol and drugs use among adolescents

Ermelinda Caldeira^{1,2}, Ana Dias², Lara Pinho^{1,2},
Anabela Coelho^{1,2}, Leonel Lusquinhos², Manuel Lopes^{1,2}

¹Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ²Nursing Department, Universidade de Évora, Évora, Portugal

Keywords: Adolescence, Health Risk Behaviors, Illicit Drugs; Licit Drugs

Objective: To understand and analyze the prevalence and factors associated with the use of tobacco, alcohol and other drugs in grade students.

Methods: Observational study included in the AlenRiscos Observatory. The sample is composed of 12767 adolescents from public and private schools of the 3rd cycle of basic education (7th and 9th grades) in the Alentejo region. Data collection was carried out during 5 school years, the last one in 2021/2022. An online self-completion questionnaire was applied from the LimeSurvey platform, accessible by the University of Évora.

Results: Most of the participants were female (52%) and aged between 11 and 18 years. 50.4 % of students were in 7th grade and 49.6% were in 9th grade. During the period under observation, the consumption of tobacco, alcohol and drugs fluctuated, with alcohol being the most consumed substance, followed by drugs and finally tobacco, in all school years. Regarding alcohol consumption, the school year with the highest consumption was 2016/2017 (48.6%). The results showed a significant decrease in consumption over the years, however, there is a higher consumption with increasing student age.

Conclusions: The evaluation of results regarding the consumption (of licit and illicit addictive substances) in adolescents allows us to define preventive measures at earlier ages, thus avoiding morbidity and mortality in adulthood.

Abstracts

Effects of a yoga practice on respiratory capacity in military pilots from the Portuguese Airforce

Sara Santos¹, Santos Villafaina², José Parraça¹,
Orlando Fernandes¹, Filipe Melo^{1,3}

¹Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ²Universidad de Extremadura; ³Laboratório de Comportamento Motor, Faculdade de Motricidade Humana, Universidade de Lisboa

Keywords: Airforce pilots; Ashtanga vinyasa yoga; Forced expiration volume; Pranayama

Objective: Spirometry and respiratory training modalities in the airforce context are of great importance. It is especially relevant in cases where pilots are deployed to aircraft flying missions at altitudes where air is rarefied. This specific training might even impact focus and mission performance in these less oxygenated environments. The goal of this study was to verify the effects of a yoga practice on respiratory capacity in military pilots from the Portuguese Airforce.

Methods: A total of 18 military pilots in the Portuguese Air Force Academy course “Masters in Military Aeronautics: aviator pilot specialist” participated in this randomized controlled trial. Respiratory capacity was analyzed with a spirometer (Vitalograph copd-6™) before and after a 12-week yoga program. Participants were randomly divided in yoga class (intervention n=10) and waiting list (control n=8); the intervention group had practice twice a week while control group did only the regular course training from the Portuguese Air Force. Data was analyzed with Jamovi (version 2.3.26), parametric tests were conducted: differences at baseline were studied using the t-test for paired samples.

Results: A pulmonary alteration was noted in FEV1% (p=0.017) for the intervention group with Mean (Standart Deviation (SD)) baseline values being 95.300 (12.0467) and becoming 120.750 (24.5459) after the 12-week yoga protocol, but not for the control group (p=0.135).

Conclusions: Military pilots already have a strong adaptative physiological response to stress due to the military training they receive. Even then, the practice of yoga leads to differences in forced expiration volume that are related to an increase in the strength and endurance of respiratory muscles following yoga training.

Functional Capacity, Quality-Of-Life and “Patient Satisfaction” with Nursing Care of Older Adults

Margarida Goes^{1,2}, Henrique Oliveira^{2,3}

¹Escola Superior de Enfermagem São João de Deus, Universidade de Évora, Portugal; ²Comprehensive Health Research Centre, Escola Nacional de Saúde Pública, Lisboa, Portugal; ³Instituto Politécnico de Beja, Portugal

Keywords: Aging; Functionality; Quality of Life; Satisfaction; Nursing Interventions

Objective: Firstly, to determine the Functional Capacity profile of people aged 65 and over residing in the Baixo Alentejo Region (BAR) based on a self-care model and standardize it concerning the age of the respondents. Second, to determine the

Quality of Life (QOL) profile of the same individuals, identifying the biological and sociodemographic predictors of the QOL domains. Third, to quantify the "Patient Satisfaction" indicator as a sensitive measure of nursing care.

Methods: Cross-sectional, descriptive study. A random sample stratified by gender and age group was used, with 351 people aged 65 and over residing in BAR. A confidence level of 95% and an error margin of 4.5% were adopted. Several statistical analysis techniques were used: psychometric, descriptive, and inferential. Mainly used were Confirmatory Factor Analysis as an application of Structural Equation Modelling, Multiple Linear Regression Methods, and Ordinal Regression Methods.

Results: The standardization of Functional Capacity profiles as a function of age revealed that up to 74 years of age, the "NO problem" profile presents the highest probability. Beyond this age, the "MILD PROBLEM" profile shows the highest probability, and after 85 years and over, the "MODERATE Problem" profile presents a higher probability in relation to the former. "Education level" and "living together" were considered the biological and sociodemographic predictors of QoL domains. The most important of all facets were "Activities of daily living", "Work capacity", "Self-esteem", and "Opportunities to acquire new information and skills". The population sample showed a "Patient Satisfaction" score of 62.1% concerning the nursing care received.

Conclusions: It is proposed to assess the needs of nursing care of older adults residing in BAR based on the assessment of their Functional Capacity, QoL, and "Patient Satisfaction".

Keratinocyte Cancer Prevention at Primary Healthcare Centres in Portugal

Maria Miguel Castella¹, Marília Silva Paulo²,
Mélanie Raimundo Maia³, Luís Velez Lapão^{2,3,4}

¹Escola Nacional de Saúde Pública, Universidade Nova de Lisboa, Portugal; ²Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal; ³UNIDEMI, NOVA School of Science and Technology, Universidade Nova de Lisboa, Portugal; ⁴LASI, Intelligent Systems Associated Laboratory, Guimarães, Portugal

Keywords: Clinical pathway; Keratinocyte cancer; Prevention, Primary health care; Health information systems; Skin cancer

Objective: To identify the clinical pathway of the patients diagnosed, or with suspicion of with keratinocyte cancer in the Agrupamento de Centros de Saúde Arco Riberinho (ACES-AR), that comprises Alcochete, Barreiro, Moita and Montijo, providing care to 219359 people.

Methods: Two focus groups were conducted. A randomized number was assigned to all the Family Physicians of the ACES, a selection of two per healthcare organization was done and they were invited to participate in this study. The selection and invitation procedures were done by the Clinical Research Group of the ACES-AR. Eleven physicians have agreed to participate, but just six, from three healthcare organizations, were able to participate. All participants have signed an informed consent and interviews were recorded.

Results: The focus groups were conducted by a trained researcher in qualitative methods and assisted by a second peer.

Researchers have used a semi-structured guide to conduct the process and achieve the goal of the study. Interviews were transcribed and analysed on NVIVO 10. Main themes emerged were: distance care (from dermatologist), hospital referral system, and skin lesions.

Conclusions: Family physicians are concerned with the topic as their district hospital does not have capacity of response in dermatology, and other hospitals are considered too far-away by patients. Patients' follow-up for this disease is minimal as Family Physicians usually lose contact with the patient's information on treatment. There is no integration of medical information systems. There is a lack of technical infrastructure to facilitate the referral triage process (such as digital cameras).

Metabolome profile of critically ill patients regarding invasive mechanical ventilation

Tiago A. H. Fonseca^{1,2}, Cristiana Von Rekowski^{1,2}, Rúben Araújo^{1,2},
Luís Bento^{2,4}, Cecília R. C. Calado^{1,5}

¹ISEL - Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, Rua Conselheiro Emídio Navarro 1, 1959-007 Lisboa, Portugal; ²Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal; ³Intensive Care Department, Centro Hospitalar Universitário de Lisboa Central, CHULC; Rua José António Serrano, 1150-199 Lisboa, Portugal;

⁴Integrated Pathophysiological Mechanisms, CHRC, NOVA Medical School, Faculdade de Ciências Médicas, NMS, FCM, Universidade NOVA de Lisboa; Campo Mártires da Pátria, 130, 1169-056 Lisboa, Portugal; ⁵CIMOSM - Centro de Investigação em Modelação e Optimização de Sistemas Multifuncionais. ISEL - Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, Rua Conselheiro Emídio Navarro 1, 1959-007 Lisboa, Portugal

Keywords: ICU; COVID-19; Metabolomics; FTIR spectroscopy; Invasive mechanic ventilation

Objective: Invasive mechanical ventilation (IMV) is used to treat critically ill patients, including those with COVID-19. In this exploratory analysis, we aimed to perform a comparative analysis of the clinical characteristics and serum metabolomic profile between ventilated and non-ventilated COVID-19 patients.

Methods: Sixteen critically ill COVID-19 patients from an intensive care unit (ICU) were included, with half being submitted to IMV. The patient groups didn't differ in age, sex, or comorbidities. Serum metabolome was obtained after macromolecules precipitation by a mixture of methanol, acetonitrile and water. The molecular profile associated to the serum metabolome was acquired by Fourier-transform infrared (FTIR) spectroscopy between 400-4000 cm⁻¹, with a 2 cm⁻¹ resolution. Spectra were pre-processed by diverse methods, including baseline correction and different kinds of normalizations, and analyzed using principal component analysis (PCA) associated or not with linear discriminant analysis (LDA).

Results: The serum metabolome acquired by FTIR spectroscopy enabled the development of a PCA-LDA model that predicted whether patients were under IMV, with 83% accuracy in the validation set, and that the serum metabolome was significantly different between the two groups. The PC2 loading vector

highlighted two relevant zones. One of them in the fingerprint zone (1028 to 1170 cm^{-1}) and another (between 1585 and 1686 cm^{-1}) that involved the vibrational frequencies of N-H and C=O bonds, such as from peptides' amides, possibly related to increased proinflammatory cytokines. This may be due to the increased inflammatory status from ventilated patients, also evidenced by increased hematological markers (e.g., C-reactive protein at $p<0.03$), and higher levels of blood glucose and lactate ($p<0.001$).

Conclusions: The molecular fingerprint associated with the serum metabolome enabled the detection of the general metabolic impact of IMV. This technique can be applied in an economic, rapid, and high-throughput mode to detect critical metabolic shifts, leading to better patient management.

Effects on Cardiovascular Risk Factors: A Randomized Controlled Trial comparing HIIT and MICT in Coronary Artery Disease Patients

Catarina Gonçalves^{1,2}, Jorge Bravo^{1,2}, João Pais³,
Armando Raimundo^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ³Hospital Espírito Santo de Évora, Évora, Portugal

Keywords: Cardiovascular Diseases; Cardiorespiratory fitness; Cardiovascular Risk Factors; High-intensity interval training; Randomized Controlled Trial

Objective: Investigate the effects of two community-based exercise programs using two short-term protocols: High-intensity interval training (HIIT) and moderate-intensity continuous training (MICT) in cardiovascular risk factors in stable coronary artery disease (CAD) patients.

Methods: We conducted a RCT, recruiting 46 CAD patients from Hospital Espírito Santo of Evora. Participants were randomized to three sessions weekly HIIT ($n = 23$) or MICT ($n = 23$) for 6 weeks. HIIT consisted of 4×4 min intervals of vigorous exercise ($\approx 85\text{--}95\%$ maximum capacity) interspersed with 1 min periods of recovery. MICT was 28 min of moderate-intensity continuous exercise ($\approx 70\text{--}75\%$ maximum capacity). The primary outcome was the change in cardiorespiratory fitness [peak oxygen uptake (VO_2peak)] after 6 weeks. Secondary outcomes included cardiovascular disease risk biomarkers, blood pressure, body composition, physical activity (PA) levels, and health-related quality of life.

Results: At 6 weeks, VO_2peak improved more with HIIT (2.3 $\text{ml kg}^{-1}\text{min}^{-1}$, $\Delta\text{m2-m1\%HIIT}$: 14%) compared with MICT (1.4 $\text{ml kg}^{-1}\text{min}^{-1}$, $\Delta\text{m2-m1\%MICT}$: 9%). HIIT also showed significantly effects on sedentarism time ($\Delta\text{m2-m1\%HIIT}$: 15% vs. $\Delta\text{m2-m1\%MICT}$: 10%), body fat mass ($\Delta\text{m2-m1\%HIIT}$: 4.5% vs. $\Delta\text{m2-m1\%MICT}$: 3.2%), waist circumference ($\Delta\text{m2-m1\%HIIT}$: 4.1% vs. $\Delta\text{m2-m1\%MICT}$: 2.5%), HbA1c ($\Delta\text{m2-m1\%HIIT}$: 10.4% vs. $\Delta\text{m2-m1\%MICT}$: 32.3%) and TSH ($\Delta\text{m2-m1\%HIIT}$: 16.5% vs. $\Delta\text{m2-m1\%MICT}$: 3.1%).

Conclusions: In stable CAD, HIIT improved cardiorespiratory fitness, fat loss, and decreased sedentarism time more than MICT by a clinically meaningful margin. HIIT is a safe, well-

tolerated, and clinically effective intervention that produces short-term improvement in cardiovascular risk factors. It should be considered by all cardiac rehabilitation programs as an adjunct or alternative to MICT.

Exercise-based Programs for cardiac patients: HIIT vs MICT – A Randomized Controlled Trial comparing Fat Loss and Blood Biomarkers

Catarina Gonçalves^{1,2}, Jorge Bravo^{1,2}, João Pais³,
Armando Raimundo^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ³Hospital Espírito Santo de Évora, Évora, Portugal

Keywords: Cardiovascular Disease; Cardiovascular Risk Factors; Clinical Trials; High Intensity Interval Training; Randomized Controlled Trial

Objective: The aim of this study was to investigate the effects of two exercise-based programs using two short-term (six-week) protocols: High-intensity interval training (HIIT) and moderate-intensity continuous training (MICT), compared to the control group in fat loss and blood biomarkers.

Methods: 72 patients with coronary artery disease (CAD) were individually randomized (1:1:1) into three groups: HIIT, MICT, and control. Both training programs consisted of 6 weeks of supervised treadmill exercise, three sessions per week. The MICT at $\approx 70\text{--}75\%$ of heart rate (HR) peak and HIIT at $\approx 85\text{--}95\%$ of HRpeak. The control group were given recommendations to adopt healthy lifestyles. Outcome measurements included an assessment of anthropometric variables (DXA) and blood biomarkers.

Results: The HIIT group demonstrated greater improvements comparing to MICT in body fat mass ($\Delta\text{m2-m1\%HIIT}$: 4.5%, $p < .001$ vs. $\Delta\text{m2-m1\%MICT}$: 3.2%, $p < .001$), waist circumference ($\Delta\text{m2-m1\%HIIT}$: 4.1%, $p = .002$ vs. $\Delta\text{m2-m1\%MICT}$: 2.5%, $p = .002$), HbA1c ($\Delta\text{m2-m1\%HIIT}$: 10.4%, $p < .001$ vs. $\Delta\text{m2-m1\%MICT}$: 32.3%, $p < .001$) and TSH ($\Delta\text{m2-m1\%HIIT}$: 16.5%, $p = .007$ vs. $\Delta\text{m2-m1\%MICT}$: 3.1%, $p = .201$). Both HIIT and MICT exercise protocols promoted a significant improvement in anthropometric variables and blood biomarkers compared to the control group.

Conclusions: This RCT showed that both training programs were equally effective in improving fat loss and blood biomarkers in CAD patients, but the HIIT group showed further improvements compared to MICT. On the contrary, not doing any type of exercise-based program after a cardiovascular problem has shown worse results.

Improving health outcomes in myocardial infarction patients with two short-term protocols:

A randomized controlled trial

Catarina Gonçalves^{1,2}, Jorge Bravo^{1,2}, João Pais³,
Armando Raimundo^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ³Hospital Espírito Santo de Évora, Évora, Portugal

Objective: Investigate the effects of two community-based exercise programs using two short-term (six-week) protocols: High-intensity interval training (HIIT) and moderate-intensity continuous training (MICT) in physical fitness and physical activity (PA) levels in myocardial infarction (MI) patients.

Methods: Physical fitness (aerobic capacity and muscle strength) and daily PA (accelerometry) were assessed before and after six weeks of intervention in 69 patients diagnosed with MI. All patients were randomly assigned to the two exercise groups (HIIT or MICT) or the control group (no exercise). Both training programs consisted of six weeks of supervised treadmill exercise, three sessions per week. The MICT at \approx 70-75% of heart rate (HR) peak and HIIT at \approx 85-95% of HRpeak. The control group only did the medical recommendations.

Results: Both HIIT and MICT could significantly improve VO_2peak , but HIIT increased considerably by 14% ($\Delta=2.3 \text{ ml kg}^{-1}\text{min}^{-1}$, $d = 1.54$) compared with 9% of MICT ($\Delta=1.4\pm1.2 \text{ ml kg}^{-1}\text{min}^{-1}$, $d=0.68$). HIIT also showed more positive effects on sedentarism time with a decrease of 15% ($\Delta= -148.6\pm106.1 \text{ min/day}$, $d = -1.20$) compared with 10% of MICT ($\Delta= -105.5\pm88.0 \text{ min/day}$, $d = -0.91$). Moreover, the control group showed negative results on physical fitness and daily PA measurements and sedentarism time.

Conclusions: HIIT can improve physical fitness and daily PA more considerably than MICT without adversely affecting patient safety. These findings indicate that HIIT may be an alternative and effective training method in community-based exercise programs for patients with MI. On the contrary, not doing any exercise-based program after a cardiovascular problem has shown worse results.

The future of digitalisation in chronic disease management in Primary Health Care: a foresight case study in Lisbon Health Region

Mariana Peyroteo^{1,2}, Mélanie Maia^{2,3}, Luís Velez Lapão¹⁻⁴

¹Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal; ²UNIDEMI, Department of Mechanical and Industrial Engineering, Faculty of Science and Technology, Universidade NOVA de Lisboa, Caparica, Portugal; ³LASI, Intelligent Systems Associated Laboratory, Guimarães, Portugal, ⁴WHO Collaborating Center on Health Workforce Policy and Planning, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Portugal

Keywords: Chronic Diseases; Digital Health; Foresight; Primary Health Care; Strategic Scenarios; Primary Health Care

Objective: Due to the pressure on healthcare systems, the COVID-19 pandemic has forced administrators to quickly leverage and integrate existing technology to provide remote consultations to reach chronically ill patients. As such, this study aims to develop scenarios about the digitalisation of PHC in the Portuguese Healthcare System to support evidence-based policymaking.

Methods: This study follows a collaborative foresight approach using a scenario development methodology. Based on previous literature reviews and stakeholder interviews, a conceptual model of PHC services was developed to consider the role of information (e.g. supported by digital services). The study included three focus groups involving primary managers and health professionals with a ten-year timeframe.

Results: The driving forces that may influence the digitalisation of PHCs over the next ten years were identified as the potential for service innovation, the role of governance and regulations, and the impact of climate change. These resulted in the design of three relevant and plausible scenarios: "Digital PHC and active patients", "NHS transition", and "Climate exigency". The scenarios where governance and climate change stand out showed limited benefits compared to the scenario where there is technology investment. In this, combining innovation skills to establish digital services with improved governance policies enabling direct investment in digital technologies would go much further towards the goals of more patient-centred care.

Conclusions: The collaborative reflection process to identify the driving forces made the different stakeholders engage, discuss the different points of view and identify essential measures to be taken in different scenarios. Furthermore, it allowed the understanding of the impact of the actions to be taken for PHC digitalisation to be implemented most effectively, enabling the sustainability of the National Health Service in the Lisbon region, which up to this point has only been discussed at a limited and theoretical level.

Pharmacogenomic biomarkers as a source of evidence for effectiveness and safety in Alzheimer's disease therapy

Ana Paula Silva¹, Margarida Perdigão²,
Margarida Espírito Santo^{1,3,4}, Ana Margarida Advinha^{2,5,6}

¹Escola Superior de Saúde, Universidade do Algarve, Faro, Portugal; ²Unidade Regional de Farmacovigilância do Centro e Norte Alentejano, Escola Superior de Enfermagem São João de Deus de Évora, Universidade de Évora, Évora, Portugal; ³Centro de Estudos e Desenvolvimento em Saúde (CESUALg), Universidade do Algarve, Faro, Portugal; ⁴ABC-RI, Algarve Biomedical Center Research Institute, Faro, Portugal; ⁵CHRC – Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ⁶Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal

Keywords: Alzheimer disease; Biomarkers; Pharmacogenomic; Safety; Effectiveness

Objective: This study aimed to describe the pharmacogenetic biomarkers, contained in the Summary of Product Characteristics (SmPC) of the medicines with marketing authorization applications in Portugal, as a source of effectiveness and safety profile for medicines used in the therapy of Alzheimer's disease.

Methods: The work was developed in two phases: i) the search for pharmacogenomic biomarkers in SmPC of the medicines used in the therapy of Alzheimer's disease with marketing authorization in Portugal; and ii) a systematic literature review based on the data obtained in the first phase, with objective of finding studies in international literature that could describe and characterize the biomarkers found before and possibly identifying other relevant biomarkers. Finally, the levels of evidence and recommendation grades were classified for every study, according to Oxford Centre for Evidence-Based Medicine classification.

Results: Were identified 4 drugs with marketing authorization in Portugal to be used in Alzheimer disease treatment, and pharmacogenomic information was pointed for all. The pharmacogenomic biomarker identified more often was CYP2D6, followed by CYP2E1, CYP3A4, CYP1A2, CYP2C9. These results will be mostly supported by the systematic review. More detailed information should be considered and included in the main medicine's information source.

Conclusions: Most pharmacogenomic variants are not studied or acknowledged by the genetic tests and still need more scientific research that confirm their usefulness. Therefore, only a small number of variants is considered when prescribing drugs used in the therapy of Alzheimer's disease. It is expected that the results and information obtained through the systematic review contribute to develop more actual recommendations for each medicine pharmacogenomic profile into the SmPC, representing an improved tool for clinical decisions.

Portuguese Global Medicines Access Index 2022: Intercalary study

Sofia de Oliveira-Martins^{1,2}, Ana Margarida Advinha^{1,3}

¹Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ²Faculdade de Farmácia da Universidade de Lisboa, Lisboa, Portugal; ³Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal

Keywords: Access to medicines, Healthcare quality indicators, Health services accessibility, Real-world evidence

Objective: To study the process of introducing new drugs in Portuguese National Health Service (NHS) hospitals, after funding decision; to understand how results are measured in hospitals; to deepen the impact of drug shortages; and to identify barriers to accessing medication.

Methods: Cross-sectional, observational study (period under analysis – 2021). Data collection was carried out through an electronic questionnaire, sent to all NHS hospitals (September 15 and October 15, 2021.) Participation was voluntary.

Results: The response rate was 45% ($n=22/N=49$). Regarding the introduction of innovative drugs, 95% of the hospitals reported having a specific internal procedure, prior to this introduction, after a funding decision. Only 68% ($n=22$) of the responding institutions stated that they select the drugs used based on their real value. However, only 27% reported performing data management in the real-world context. Half of the respondents stated that shortages constituted a serious problem affecting all types of medicines. Only 27% had implemented pharmaceutical consultations, a fact that is worrisome, having attention to the need to control ambulatory patients. The main barriers identified to access to medicines were administrative burden (68%), lack of human resources (55%) and medication financing model (41%).

Conclusions: In this study, previously identified barriers to access to medicines were studied, namely the payment model, lack of human resources, acquisition process, and administrative burden. Results point to the need to rethink the health system organization at the hospital level, to allow early access to innovation.

How is low back pain being managed in primary healthcare? Findings from a scoping review on models of care

Susana Tinoco Duarte^{1,2}, Alexandre Moniz^{3,4}, Daniela Costa^{3,4}, Bruno Heleno³, Pedro Aguiar^{1,2}, Eduardo Brazete Cruz^{5,6}

¹Comprehensive Health Research Center, Escola Nacional de Saúde Pública, Universidade NOVA de Lisboa, Lisbon, Portugal;

²Escola Nacional de Saúde Pública, Universidade NOVA de Lisboa, Lisbon, Portugal; ³Comprehensive Health Research Center, NOVA Medical School | Faculdade de Ciências Médicas, Universidade NOVA de Lisboa, Lisbon, Portugal; ⁴EpiDoc Unit, NOVA Medical School | Faculdade de Ciências Médicas, NMS|FCM, Universidade Nova de Lisboa; Lisbon, Portugal;

⁵Comprehensive Health Research Center, Universidade NOVA de Lisboa, Lisbon, Portugal; ⁶Physical Therapy Department, Polytechnic Institute of Setúbal, Setúbal, Portugal

Keywords: Implementation science, Low back pain, Models of care, Primary care

Objective: Models of care (MoCs) are frameworks that describe evidence-informed healthcare that should be delivered to patients in a given setting. This study aimed to synthesize research evidence regarding the core characteristics, context features and key common elements of MoCs implemented in primary healthcare for the management of low back pain (LBP).

Methods: Searches on MEDLINE(Pubmed), EMBASE, Cochrane Central Register of Controlled Trials, PEDro, Scopus, Web of Science and grey literature databases were conducted. Handsearching in relevant peer-reviewed journals and organisations' websites was also performed. Title/abstracts and full-text screenings were carried out independently by two researchers and disagreements were solved through consensus. Eligible records report findings of MoCs implemented for LBP patients in primary healthcare settings, which were investigated through a descriptive qualitative content analysis.

Results: Thirty studies reporting 12 MoCs were included. The majority evaluated acceptability, implementation strategies, clinical effectiveness and cost-effectiveness of new MoCs, mostly through randomized controlled trials and qualitative studies. All MoCs were implemented in high-income countries and had clear objectives, which included, but were not limited to, delivery of guideline-concordant care and reduction of healthcare resources utilization. In general, most MoCs followed a stratified care approach. Assessment of LBP patients typically occurred in primary healthcare, while the care delivery usually took place in community-based settings or outpatient clinics. The involvement of general practitioners and physiotherapists was reported in the majority of the studies. Education and exercise were the most common health interventions, but intervention content, follow-ups and discharge criteria were not fully reported.

Conclusions: MoCs for LBP in primary healthcare share relevant key elements, but report of interventions is heterogeneous. This study provides a comprehensive understanding of the characteristics of the existing MoCs, which is essential new information to underpin future research implementation studies for LBP.

Models of care for low back pain patients in primary healthcare: a scoping review on implementation processes and outcomes

Susana Tinoco Duarte^{1,2}, Alexandre Moniz^{3,4}, Daniela Costa^{3,4}, Bruno Heleno³, Pedro Aguiar^{1,2}, Eduardo Brazete Cruz^{5,6}

¹Comprehensive Health Research Center, Escola Nacional de Saúde Pública, Universidade NOVA de Lisboa, Lisbon, Portugal;

²Escola Nacional de Saúde Pública, Universidade NOVA de Lisboa, Lisbon, Portugal; ³Comprehensive Health Research Center, NOVA Medical School | Faculdade de Ciências Médicas, Universidade NOVA de Lisboa, Lisbon, Portugal; ⁴EpiDoc Unit, NOVA Medical School | Faculdade de Ciências Médicas, NMS|FCM, Universidade Nova de Lisboa; Lisbon, Portugal;

⁵Comprehensive Health Research Center, Universidade NOVA de Lisboa, Lisbon, Portugal; ⁶Physical Therapy Department, Polytechnic Institute of Setúbal, Setúbal, Portugal

Keywords: Implementation science, Low back pain, Models of care, Primary care

Objective: Models of care (MoCs) are frameworks that describe evidence-informed healthcare that should be delivered to patients in a given setting. This study aimed to map the current literature about the implementation of MoCs for low back pain (LBP) in primary healthcare, analysing their implementation strategies, process evaluations and changes in patient-, organizational- and implementation-level outcomes.

Methods: This study followed the Joanna Briggs Institute guidelines for scoping reviews. Searches on MEDLINE(Pubmed), EMBASE, Cochrane Central Register of Controlled Trials, PEDro, Scopus, Web of Science and grey literature databases were conducted. Additionally, handsearching was also performed in relevant peer-reviewed journals and organisations' websites. Title/abstracts and full-text screenings were carried out independently by two researchers and disagreements were solved through consensus. Data concerning frameworks and implementation strategies, barriers and facilitators, outcomes, outcome measures and main results were extracted and examined through a descriptive qualitative content analysis.

Results: Thirty studies reporting 11 MoCs were included. Only 5 MoCs reported the use of frameworks/theories to develop or evaluate the implementation process. When described, implementation strategies varied across studies. The most used strategy was the training of health professionals involved in providing care. Patient-level outcomes (symptoms and disease severity, psychosocial and work-related) were reported in 11 MoCs and organizational-level outcomes (healthcare resources utilization, costs and quality assessment) were evaluated in 10 MoCs. Few studies reported implementation-level outcomes, such as acceptability, feasibility and adoption.

Conclusions: As most MoCs were not informed by implementation frameworks, the results of this scoping review are inconclusive. Systematic approaches for the development, evaluation and report of implementation processes of the MoCs are crucial to justify their effectiveness and changes in outcomes.

CkeckAP: Psoriatic Arthritis Prevalence Among Portuguese Psoriasis Patients and Evaluation of the EARP Questionnaire Performance

Ana Henriques¹, Fernando Santos², João Sousa², Laura Gago², Margarida Rocha², Mariana Santos², Ana Chícara², Anabela Barcelos^{1,3} Manuela Loureiro³, Susana Silva³, Elsa Sousa⁴, Carolina Matos⁴, Joana Ferro Antunes⁴, Miguel Alpalhão⁴, Sofia Magina⁵, Alexandra Bernardo⁵, Maria Rato⁵, Pedro Ponte⁶, Tiago Torres⁷, Tiago Meirinhos⁷, Martinha Henrique⁸, Marília Rodrigues⁸, Diogo Jesus⁸, Alexandra Daniel⁸, Luísa Brites⁸, Pedro Bastos⁹, Patrícia Nero⁹, Maria Valada¹, David Lopes¹, Rute Sousa¹, Helena Canhão^{1†}, Ana Rodrigues^{1,6}

¹CHRC, NOVA Medical School, NMS, Universidade NOVA de Lisboa; Lisboa, Portugal; ²Hospital Egas Moniz, Centro Hospitalar Lisboa Ocidental, CHLO- E.P.E, Lisboa, Portugal; ³Serviço de Dermatologia e Serviço de Reumatologia do Centro Hospitalar do Baixo Vouga, Aveiro, Portugal; ⁴Hospital de Santa Maria, Centro Hospitalar e Universitário de Lisboa Norte, Centro Académico de Medicina de Lisboa, Lisboa, Portugal; ⁵Centro Hospitalar Universitário de São João, Porto, Portugal; ⁶Hospital dos Lusíadas, Lisboa, Portugal; ⁷Instituto Médico de Estudos Imunológicos, Porto, Portugal; ⁸Hospital de Santo André, Centro Hospitalar de Leiria, E.P.E, Leiria, Portugal; ⁹Hospital CUF Descobertas, S.A, Lisboa, Portugal

Keywords: Psoriasis; Psoriatic arthritis; EARP Questionnaire

Objective: The present study aims to estimate the percentage of Psoriatic Arthritis (PsA) in psoriasis patients in the Portuguese dermatology setting. Additionally, it aims to evaluate the performance of the EARP questionnaire in Portuguese psoriasis patients and determine the appropriate cut-off point for early diagnosis of PsA.

Methods: A multicentre, cross-sectional, observational study with two independent assessments: one by the dermatologist and the other by the rheumatologist. Patients were recruited at one of eight Portuguese study sites (private and public settings). PsA diagnosis was defined by a combination of expert opinion with CASPAR criteria among psoriasis patients in a dermatology consultation setting. PsA prevalence estimates were computed as proportions together with the corresponding 95% confidence interval. Prevalence by gender and age strata was computed. PsA screening results, mediated by the EARP questionnaire, were compared with the gold standard (PsA diagnosis made by a rheumatologist) and sensitivity, specificity, positive predictive value, negative predictive value, and area under the curve were determined.

Results: 172 patients were included with a mean age (SD) of 53.8 (14.5) years, 53.5% male. The prevalence of psoriatic arthritis in patients with psoriasis in our sample was 8.70% (95% CI: 4.8-14.2). Psoriatic arthritis is more frequent in women (9.09%) than in men (8.33%) and in the 31-59 years age group. The EARP questionnaire showed a good internal consistency (Cronbach's $\alpha=0.81$). For an EARP questionnaire cut-off point of 3, the questionnaire's sensitivity and specificity were 71.4% and 40.1%, respectively. The AUC was 0.558 (95% CI: 0.429-0.687).

Conclusions: The Portuguese version of the EARP questionnaire has an acceptable sensitivity, albeit with a low specificity.

Nevertheless, the Portuguese version of the EARP (EARP-PT) questionnaire is a helpful tool for the early detection of PsA symptoms by dermatologists when assessing patients who are susceptible to this condition.

Patient-reported outcome measures in cardiac rehabilitation programs: preliminary results of a Systematic Review

José Moreira¹, Duarte Tavares², Armando Raimundo³, Bruno Delgado⁴, Paulo Boto¹

¹Comprehensive Health Research Centre, National School of Public Health, Lisbon, Portugal; ²Lisbon Metropolitan Transports, Lisbon, Portugal; ³Comprehensive Health Research Centre, University of Évora, Évora, Portugal; ⁴Catholic University of Portugal, Institute of Health Sciences, Porto, Portugal

Keywords: Cardiac rehabilitation; Patient-reported outcome measures; Quality of life; Randomized controlled trial; Treatment outcome

Objective: Patient-reported outcome measures (PROMs) are recommended as important endpoints in cardiac rehabilitation programs (CRP), highlighting patients' health status, functioning and health-related quality of life (HrQol), as well as enabling clinical and health decisions to be made in an informed manner. Analyze which PROMs are used to assess the HrQol of patients participating in a CRP following a cardiac event, as a function of assessment time.

Methods: A systematic review of peer-reviewed English language articles was undertaken according to recommendations of the PRISMA group, conducted in MEDLINE, CINAHL and Cochrane Library from date of inception up to January 2017 and up to December 2022. The PICO strategy was used, to select RCTs studies comparing CRP against usual care, focusing time on HrQol assessment through PROMs. The protocol registered with PROSPERO: CRD42022344240.

Results: A total of 533 articles were found and 453 articles were eliminated after reading the title, abstract and removing duplicates. After reading full text the sample was 13 articles. The total number of participants was 4194, 2858 male, with a mean age of 61.7 years, and conducted in different countries: Germany and USA[n=2], Italy, UK, Lithuania, Egypt, China, Israel, Pakistan, UAE, Netherlands[n=1]. The intervention included the four phases of a CRP, using different PROMs to assessment HrQol (EQ5D[n=4], SF36[n=2], SF12[n=3], KCCQ, HeartQol, MacNewQLMI[n=1]) and different moments since the end of CRP over a period quantified in months: 3[n=6], 6[n=5], 12[n=6], 1, 8, 9, 18, 24, 36, 48 months[n=1].

Conclusions: Although different PROMs in CRP, the most prevalent HrQol assessment time was at 3, 6 and 12 months after program start.

Effects of an eight-weeks High-Speed Resistance Training program on heart rate variability in community-dwelling independent older adults

Alexandre Duarte Martins^{1,3}, Orlando Fernandes¹,
João Paulo Brito^{2,3,4}, Nuno Batalha¹

¹Department of Sport and Health, School of Health and Human Development, University of Évora, 7004-516 Évora, Portugal;

²Comprehensive Health Research Centre (CHRC), University of Évora, Portugal; ³Life Quality Research Centre, Rio Maior 2040-413, Portugal; ⁴Research Centre in Sport Sciences, Health Sciences and Human Development, Vila Real 5001-801, Portugal

Keywords: Older adults; Resistance training; Velocity-based training; Variability; Heart Rate

Objective: This study examined the effects of an eight-week high-speed resistance training (HSRT) program on heart rate variability (HRV) indices (time and frequency domain).

Methods: This study included thirty-nine physically independent older adults (age, 68.50 ± 3.52 years; BMI, 27.88 ± 4.37 kg/m²). The HSRT program lasted eight weeks, with three sessions per week with 50–60min, each session being comprised of five-six exercises, two-three sets, and six-ten reps/exercise. The intensity was gradually increased after each session in accordance with the movement velocity (>1.3 to 0.75 m/s) representing approximately 20% to 60% of one repetition maximum. Participants executed the exercises rapidly and explosively making all repetitions for each shortening phase (concentric phase) performed as quickly as possible with the lengthening phase of the muscle (eccentric phase) being controlled for 2–3 s. The velocity of the concentric phase in each exercise was monitored through a BEAST™ sensor (Beast Technologies, Brescia, Italy). The heart rate band has been used to assess the variation in sinus origin heartbeat intervals (Polar® H10, Kempele, Finland). Using the Kubios HRV software (Kubios HRV, University of Kuopio, Finland), HRV indices were calculated. The Ethics Committee of the University of Évora approved this study (approval no. 22030). A paired sample t-test (with a cut-off of $p < 0.05$ for statistical significance) was used for comparing each outcome.

Results: After the intervention period, the HRV indices showed significant improvements: on SDNN (Δ change=22.42; $p=0.028$; effect size [ES]=0.206); RMSSD (Δ change=28.54; $p=0.022$; ES=0.213); pNN50% (Δ change=52.25; $p=0.045$; ES=0.238); and on HF (Δ change=48.65; $p=0.023$; ES=0.150). There were no differences on Average RR (Δ change=-0.79; $p=0.615$; ES=-0.055); LF (Δ change=-0.44; $p=0.974$; ES=0.01); and on the Ratio LF/HF (Δ change=-1.98; $p=0.899$; ES=-0.023).

Conclusions: The results suggest that the HSRT program is an effective and safe exercise approach to improve several HRV indices, especially SDNN and RMSSD indices.

Analysis of postural control through linear and non-linear methods throughout aging

Carolina Cabo^{1,2}, José Parraça^{1,2}, Orlando Fernandes^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano (Universidade de Évora, Évora, Portugal); ²Comprehensive Health Research Centre (CHRC) (Universidade de Évora, Évora, Portugal)

Keywords: Aging; Balance; Falls; Force platform; Postural control

Objective: The natural aging process is defined as continuous and irreversible and is associated with a decline in physical and cognitive functions. The maintenance of balance and body orientation in humans is guaranteed by the proper functioning of the Postural Control system. This study evaluated the effect of sensory-motor training on postural control in pre-retirement adults. The study evaluated the effectiveness of an active retirement program on postural control over six months of intervention, through evaluating the relative sensitivity of center of pressure measures to age-related changes in postural stability.

Methods: Measures of postural stability in the time and frequency domain were compared between a group of nine elderly adults (≤ 64 years), a group of fifty-nine elderly (65–74 years), and a group of thirty-eight elderly (≥ 75 years) in the eyes open and eyes closed conditions.

Results: The measurements identified statistically significant differences in the group ≤ 64 years in the amplitude of displacement and approximate entropy in the anteroposterior position with eyes open and in the mediolateral position with eyes closed. In the age group of 65–74 years, there were no significant differences in variables from linear methods, however there were differences between groups in variables of approximate entropy and scaled correlation. In the ≥ 75 years group, the tests performed revealed that there are statistically significant differences between the groups in the variables of total excursion and mean velocity and that there are statistically significant differences between the groups in all the variables of approximate entropy and dimensional correlation with the exception of anteroposterior position with eyes closed. In addition, there are significant differences in the comparison within the group in the three age groups evaluated.

Conclusions: In conclusion, both groups showed significant improvement in most of the analyzed variables; however, the improvements in the exercise group were significantly greater.

A view on bullying by Focal Groups in a Group of Schools in Central Alentejo

Hermenegildo Pateiro^{1,2,3}, Lauréncia Gemito^{1,3}, Ana Rita Pedro^{1,3,4}

¹Comprehensive Health Research Centre; ²Escola Nacional de Saúde Pública, Lisboa, Portugal; ³Universidade de Évora, Évora, Portugal; ⁴New Medical School, Lisboa, Portugal

Keywords: Bully; Bullying; Literacy; Perception; Victim

Objective: To characterise the way in which the phenomenon of bullying is perceived by adults in the educational community.

Methods: Focus group with analysis of responses through categories. Two focus groups were created with homogeneous constitution between them. Each group was composed of 10 individuals with the same professional characteristics. However, each group had heterogeneous characteristics in its constitution, including teachers, operational assistants, specialised technicians, managers and parents. With the organization of focus groups we intended to verify the perception of individuals about the phenomenon of bullying, as well as the perception of it by bullies and victims. We also tried to characterise the adults responsible for the students involved.

Results: After the analysis we can conclude that the perception of bullying is essentially referred to as a phenomenon repeated over time that causes real distress to the victim. This repeated process is extremely facilitated by the systematic use of social media. It was also concluded that both victims and bullies have problems at the level of social interaction. Bullies are often unaware of the harm they are doing to the victim. Victims are often fragile and easy targets for the bully, who is often part of people the victim trusts. Families sometimes initially devalue the phenomenon, and sometimes, by the time they appreciate it, it's too late to try to solve it. It was also found that the participants think it is very important to have a comprehensive intervention plan to prevent this phenomenon.

Conclusions: The phenomenon of bullying is increasingly serious and causes lifelong damage to both the bully and the victim. The social media helps to increase the cyber bullying, because the bully is protected by your computer, can use a false identity. It is essential to prevent this phenomenon in a school environment. Focus group participants were unanimous in stating that increased literacy of parents and students can help prevent bullying.

Models of Care for People with Knee Osteoarthritis in Primary Healthcare: A Scoping Models of Care for People with Knee Osteoarthritis in Primary Healthcare: A Scoping Review

Daniela Costa^{1,2,4}, Luís A Gomes^{2,3,4}, Ana M Rodrigues^{2,4}, Daniela Gonçalves-Bradley⁵, Helena Donato⁶, Eduardo B Cruz^{2,3}

¹NOVA National School of Public Health, Public Health Research Centre, Universidade NOVA de Lisboa, Lisbon, Portugal;

²Comprehensive Health Research Center (CHRC), NOVA Medical School, Lisbon, Portugal; ³Physical Therapy Department, School of Health, Polytechnic Institute of Setúbal, Setúbal, Portugal;

⁴EpiDoC Unit, NOVA Medical School, Universidade Nova de

Lisboa, Lisbon, Portugal; ⁵Nuffield Department of Population Health, University of Oxford, Oxford, UK; ⁶Documentation and Scientific Information Service, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal

Keywords: Implementation Science; Models of care; Osteoarthritis; Primary Care

Objective: To identify Models of Care (MoCs) for knee osteoarthritis implemented in primary healthcare and describe its characteristics namely, the intervention components, the implementation and context features.

Abstracts

Methods: This scoping review, following Joanna Briggs Institute methodology, identified studies in peer-reviewed and gray literature databases. Studies about the implementation of MoCs, defined as complex interventions with an evidence-informed framework that defines the principles of care and how care should be delivered, according to local context, were included.

Results: Thirteen MoCs, among thirty studies, were included. The main objectives of these studies addressed clinical effectiveness and cost-effectiveness, evaluation of the implementation strategies using observational cohort studies, cluster RCTs or mixed-methods studies. Care was delivered mostly through stepped care pathways, but the progression, discharge criteria and follow-up plans were often poorly described. Education and structured exercise (supervised or home-based) were the most considered interventions. Physiotherapists and general practitioners were the professionals most included, but nurses, rheumatologists and pharmacists were considered, depending on context. Professionals training was often described. Some MoCs were implemented only on primary care, others also integrated community services, hospital-based and/or outpatient clinics in the care pathway. Coordination of care elements reported relied on tools to exchange information. All MoCs were implemented in high-income countries, but the context features were often poorly described. The findings of these studies suggest benefits in pain, physical function, physical activity level, healthcare utilization, quality of care, and in implementation outcomes, like fidelity and adherence.

Conclusions: This scoping review gives a comprehensive view of MoCs for knee osteoarthritis in primary care. These have been designed according to national system organization and underlying health policy, however the report of MoCs is heterogeneous. Research is needed regarding the effectiveness and process evaluation, as well as guidelines for the development and implementation report of MoCs.

Exposure of the Portuguese population to mycotoxins – the mixtures approach

Carla Martins¹, Marthe De Boevre², Duarte Torres^{3,4}, Paula Alvito⁵, Carla Nunes¹, Ricardo Assunção⁶

¹NOVA National School of Public Health, Public Health Research Centre, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Lisbon, Portugal; ²Centre of Excellence in

Mycotoxicology and Public Health, Faculty of Pharmaceutical Sciences, Ghent University, Belgium; ³Faculty of Nutrition and

Food Sciences, University of Porto, Portugal; ⁴Epidemiology

Research Unit, Institute of Public Health, University of Porto, Portugal; ⁵National Institute of Health Dr. Ricardo Jorge, Lisboa, Portugal; ⁶Egas Moniz Center for Interdisciplinary Research (CiiEM), Egas Moniz School of Health and Science, Caparica, Portugal

Keywords: Mycotoxins; Mixtures; Human biomonitoring

Objective: Mycotoxins are a group of food contaminants with associated health outcomes such as estrogenic, immunotoxic, nephrotoxic and teratogenic effects. Human biomonitoring (HBM) studies have been developed to assess the exposure to mycotoxins at the individual level and to several compounds

simultaneously. This study aimed to obtain data through a human biomonitoring study to assess the simultaneous exposure of the Portuguese population to multiple mycotoxins.

Methods: In the scope of the National Food, Nutrition, and Physical Activity Survey of the Portuguese General Population (2015-2016), 24h-urine samples from 94 participants were analyzed by liquid chromatography-mass spectrometry (LC-MS/MS) for the simultaneous determination of 42 mycotoxins' urinary biomarkers.

Results: Results revealed the exposure of Portuguese population to aflatoxins, zearalenone, deoxynivalenol, ochratoxin A, alternariol, citrinin and fumonisins B1. The human exposure to several mycotoxins was confirmed, with participants being exposed simultaneously to two (45%), three (22%), four (6%) and five (2%) mycotoxins. The most frequent combination was the binary mixture of zearalenone and deoxynivalenol. Zearalenone and alternariol, two mycotoxins with endocrine disruptive activity, were found simultaneously in 13% of participants.

Conclusions: The present study generated, for the first time and within an HBM study, reliable data on internal exposure to multiple mycotoxins at the individual level for the Portuguese population. These data are a first step for estimating the risk associated with exposure to mixtures of mycotoxins and contribute to support risk managers in the establishment of preventive policy measures to ensure public health protection.

Atmospheric levels of PM_{2.5} – the exceedance of air quality guidelines in Portugal

Lorena Lima¹, Susana Viegas², Carla Martins²

¹NOVA National School of Public Health, Public Health Research Centre, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Lisbon, Portugal; ²NOVA National School of Public Health, Public Health Research Centre, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Lisbon, Portugal

Keywords: Air quality; Particulate matter; Public health

Objective: Air pollution is the main environmental cause of mortality and premature death worldwide. The main objectives were to present the levels of exposure to PM_{2.5} in Portugal during the year 2021 and to determine the exceedance of World Health Organization air quality guidelines (WHO AQG) and EU Air Quality Directive.

Methods: Data available in the platform QualAR Portugal were collected for the year 2021. For each monitoring station, the variables collected were region, city, station, emission influence (traffic, background, industrial), and environment (urban, suburban, rural). The percentage of hourly exceedances of EU Air Quality Directive (25 µg/m³) and WHO AQG of 2006 and 2021 (10 µg/m³ and 5 µg/m³, respectively) was determined.

Results: For 2021, mean and median atmospheric levels of PM_{2.5} in Portugal were 6.53 and 4.29 µg/m³, respectively. The mean atmospheric levels of PM_{2.5} were lower when compared to the previous three-year period, except for the Center region. The percentage of hourly exceedance of guideline levels ranged from 0.07% to 11.70%, from 0.3% to 37%, and from 6% to 72%, when

compared to the EU Air Quality Directive, WHO AQG 2006 and WHO AQG 2021, respectively.

Conclusions: Despite the efforts of environmental policies aimed at increasing air quality, Portugal registered at some time points levels above the EU Air Quality Directive, being more challenging to keep within the WHO Air Quality Guidelines. Results presented herewith showed the need of continuous efforts and will support decision-making for the reduction of admissible levels of atmospheric pollutants in Portugal.

Depression and anxiety and their associated factors in university students

Pedro Amaro^{1,2}, César Fonseca^{1,3}, Anabela Afonso^{6,7}, Gonçalo Jacinto^{6,7}, Anabela Pereira^{4,5}, Lara Guedes de Pinho^{1,3}

¹Comprehensive Health Research Centre (CHRC), University of Évora, 7000-811 Évora, Portugal; ²University of Evora Institute for Advanced Studies and Research, 7004-516 Évora, Portugal;

³Nursing Department, University of Évora, 7000-811 Évora, Portugal; ⁴Centre for Research in Education and Psychology (CIEP-EU), 4200-135 Porto, Portugal; ⁵Williams James Center for Research (WJCR), 1149-041 Lisboa, Portugal; ⁶Department of Psychology, University of Évora, 7000-811 Évora, Portugal;

⁷CIMA, IIFA, University of Évora, 7000-671 Évora, Portugal;

⁷Department of Mathematics, ECT, University of Évora, 7000-671 Évora, Portugal

Keywords: Mental health; Depression; Anxiety; COVID-19; Students

Objective: To evaluate the prevalence of depressive and anxious symptoms in university students, and to compare mental health with sociodemographic characteristics.

Methods: Cross-sectional study with a sample of 3399 university students from 7 Portuguese higher education institutions was conducted. The following questionnaires were applied: sociodemographic data, Mental Health Inventory (MHI-5), Generalized Anxiety Disorder Assessment Scale (GAD-7) and Patient Health Questionnaire (PHQ-9).

Results: 75% of the participants presented mild to severe anxious symptoms and 61.2% mild to severe depressive symptoms. 6.8% of participants reported that their mental health has worsened with the COVID-19 pandemic. 19.5% reported being diagnosed with a mental disorder and of these, 38.7% reported being diagnosed after the start of the pandemic. 23% of participants reported taking medication for anxiety, depression, or another psychic problem and 11.7% reported having had thoughts that they would be better off dead or hurting themselves. The results showed better mental health for men, for students with better socio-economic status, for those who travel home more frequently and for those in a loving relationship. Students with previously diagnosed mental disorder have worse mental health.

Conclusions: There was a decrease in the mental health of university students after the pandemic compared to pre-pandemic studies and the proportion of students with anxious and depressive symptoms was alarming. It is necessary to review and readapt programs to promote the mental health of university students with the aim of improving their well-being.

An optimized method for the preliminary detection of Microplastics in the Indoor Environment

João Vasco Valente^{1,2}, Ana C. A. Sousa^{2,3,4},
M. Ramiro Pastorinho^{2,4,5}

¹Departamento de Química e CICECO, Universidade de Aveiro, Aveiro, Portugal; ²Unidade de Saúde e Ambiente, Faculdade de Ciências da Saúde, Universidade da Beira Interior, Covilhã, Portugal; ³Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ⁴Departamento de Biologia, Universidade de Évora, Évora, Portugal; ⁵Departamento de Ciências Médicas e da Saúde, Universidade de Évora, Évora, Portugal

Keywords: Confocal microscopy; Density separation; Exposome; House dust, Fluorescence microscopy

Objective: Microplastics (MPs) are plastic particles with sizes between 100 nm and 5 mm, regarded as emerging contaminants and responsible for multiple deleterious effects on human health. Due to MPs ubiquity, humans are continuously exposed. However, exposure dosimetry is very rarely performed, particularly for the indoor environment. This work describes a simple protocol to evaluate the presence of MPs in the indoor environment.

Methods: Passive dust samples were collected in two divisions (kitchen and bedroom) of the same house during one-week periods. Active dust samples were retrieved from the vacuum cleaner bag. Different methods for the extraction of MPs were tested, including the combination of density separation with sodium chloride and digestion with hydrogen peroxide. Since no standard protocol for MPs quantification in dust samples was available, the amount of dust to be used in the analysis was also optimized. The MPs were then visualized and identified by confocal microscopy after Nile Red staining.

Results: For passive samples, only the oxidative digestion of the lipids content with H₂O₂ is necessary. For active samples, it is necessary to perform a density separation with NaCl prior to oxidative digestion. Overall, the results from the passive dust samples revealed that the bedroom exhibited a higher number of fibers than the kitchen and that the highest fluorescence intensity was also registered in the bedroom samples. Concerning the active samples, the optimization of the amount of dust to be analyzed indicated that a minimum of 0.5 g of the 63 µm dust fraction should be used.

Conclusions: Nile Red visualization technique proved to be efficient in MPs detection and quantification. However, it cannot discern the composition of different MPs present in a sample. Hence, this technique should be used as a preliminary approach to confirm the existence/non-existence of MPs.

Effects of multimodal training with augmented reality in global cognition, executive function, and inhibition in older adults

Soraia Ferreira^{1,2}, José Marmeleira^{1,2}, Jesus Del Pozo-Cruz^{3,4}, Nilton Leite^{1,2}, Hélio Yoshida⁵, Armando Raimundo^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ³Departamento de Educación Física y Deporte, Universidad de Sevilla, Spain; ⁴Epidemiology of Physical Activity and Fitness Across Lifespan Research Group (EPAFit), Universidad de Sevilla, Spain; ⁵Faculdade de Educação Física (FEF), Universidade Estadual de Campinas (UNICAMP) - Grupo de Estudos em Psicologia do Esporte e Neurociências (Gepen) – Campinas (SP), Brasil

Keywords: Elderly, Multimodal training, Augmented reality training, Cognition

Objective: This study aimed to examine the effects of a multimodal intervention with augmented reality in executive functions, global cognition, and inhibition in community-dwelling elderly.

Methods: Sixty-four elderly people (72.4±5.9 years) living in the community participated in this study. Participants were divided into three groups: a control group (CG) with 22 participants, an experimental group with multimodal training (EGMT) with 21 participants, and an experimental group with multimodal training and augmented reality (EGAR) with 21 participants. All participants were assessed at the beginning of the program and after 12 weeks (at the end of the program). The experimental group exercised three times per week for 60 minutes while the control group continued their activities of daily living. In addition, global cognition was assessed with the Mini Mental State Examination (MMSE), executive functions with the Frontal Assessment Battery (FAB), and inhibition with the Stroop Test.

Results: There were significant differences when we compare the pretest and posttest within each group. In EGAR, significant differences were observed in MMSE ($p=0.020$; $\Delta\% = 2.85\%$), FAB ($p=0.013$; $\Delta\% = 6.16\%$) and ST -Color ($p=0.028$; $\Delta\% = 11.09\%$), and in EGMT in FAB ($p=0.012$; $\Delta\% = 5.52\%$). Independent samples tests showed in the changed scores, significant differences between the three groups in MMSE ($p=0.026$), but there are no differences in Stroop Test (Stroop words, $p=0.756$; Stroop color, $p=0.186$; Stroop word/color, $p=0.470$).

Conclusions: Overall, the results show that both interventions have positive effects on some of the variables studied. However, when comparing pretest and posttest within each group, EGAR has effects on more variables (MMSE, FAB and on Stroop Test -Color) than an intervention with multimodal training only (FAB).

The relationship between nursing practice environment and pressure ulcer care quality in Portugal's long-term care units

Kátia Furtado^{1,2}, Jaco Voorham³, Paulo Infante^{4,5},
Anabela Afonso^{4,5}, Manuel Lopes^{2,6}

¹Outpatient Department, Hospital of Portalegre, Unidade Local de Saúde do Norte Alentejano, 7300-312 Portalegre, Portugal; ²Comprehensive Health Research Centre (CHRC), Universidade de Évora, 7000-671 Évora Portugal; ³DTIRS—Data to Insights Research Solutions, 1750-307 Lisboa, Portugal;

⁴CIMA, IIFA, Universidade de Évora, 7000-671 Évora, Portugal;

⁵Departamento de Matemática, ECT, Universidade de Évora, 7000-671 Évora, Portugal; ⁶São João de Deus School of Nursing, Universidade de Évora, 7000-671 Évora, Portugal

Keywords: Healing; Nursing homes; Nursing Work-Revised Scale; Pressure injury; Prevention

Objective: Pressure ulcers (PUs) are responsible for high morbidity and mortality. This study aims to describe the nursing work environment in Portuguese long-term care units and to assess how this environment relates to the quality of PU care, in long term care units of Alentejo.

Methods: We looked at associations between the nursing work environment through the Nursing Work Index-Revised Scale (NWI-R) and wound healing rates obtained from a prevalence study.

Results: A total of 165 nurses completed the NWI-R-PT. Most were women (74.6%), and less than half (38.4%) had education in wound care. Of the 88 patients identified with PUs, only 63 had their PU documented. The results showed that the level of concordance with Q28 "Floating so that staffing is equalised among units" is strongly associated with a shorter PU healing time. We found no evidence for possible associations with questions on participation in policy decisions, salary level, staffing educational development and PUs healing time.

Conclusions: Good distribution of nursing staff over the units will likely improve the quality of wound care. We found no evidence for possible associations with questions on participation in policy decisions, salary level, staffing educational development and PUs healing times.

Effects of Feel-Own-Move program on the experience of the body, sleep quality and physical activity of female victims of violence. A mixed-methods study

Joana Machorrinho^{1,2}, Graça Duarte Santos^{1,2}, Guida Veiga^{1,2},
José Marmeleira^{1,2}

¹Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ²Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Évora, Portugal

Keywords: Embodiment; Psychomotor therapy; Trauma; Women; Health

Objective: Women victims of intimate partner violence (IPV) tend to experience their bodies as heavy, slow or anxious. As a result of trauma, they also feel disconnected from their body. To promote healthy bodily experiences is both a therapeutic and a social need in the field of IPV. The aim of this study was to explore the effects of Feel-Own-Move (FOM) on the experience of the body, physical activity (PA), sedentary behavior (SB) and sleep quality of female victims of IPV, through a mixed methods approach.

Methods: Nine women living in shelter homes (mean age 45.9 years; range= 32- 64 years) received FOM, an 8-week psychomotor intervention with 2 individual sessions and 1 group session per week. At week 1 (T1, baseline), week 5 (T2, pre-intervention) and week 9 (T3, post-intervention), participants were asked to use an accelerometer for seven days and nights to assess PA, SB and sleep quality. At T3, brief semi-structured interviews were conducted with each participant to explore their experience of the body and perceived internal changes.

Results: There was a decrease in SB ($\Delta\% = -16\%$) and in sleep fragmentation index ($\Delta\% = -27\%$) between T2 and T3, as well as an increase in light PA ($\Delta\% = +19\%$) although not statistically significant. Women shared experiences of increased bodily awareness regarding breathing sensations, the body-mind connection, tension-relaxation differences, and feelings of strength. Also, improvements in self-regulation, self-confidence, inner strength, motivation and vitality were reported, along with relaxation skills.

Conclusions: Feel-Own-Move showed promising results in improving bodily experiences, sleep quality and active behavior of women victims of IPV.

Gender Differences in Adolescents' Self-Knowledge Organization: The Moderating Role of Adolescence Phase

Maria João Carapeto^{1,2}, Guillem Feixas^{3,4}

¹Department of Psychology, School of Social Sciences, Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal; ³Department of Clinical Psychology and Psychobiology, Universitat de Barcelona, Barcelona, Spain; ⁴The Institute of Neurosciences, Universitat de Barcelona, Barcelona, Spain

Keywords: Implicative dilemmas; Self-concept; Self-esteem; Identification; Adolescent development; Personal constructs

Objective: To examine the effect of gender on adolescents' organization of self-knowledge and the moderation effect of adolescence phase.

Methods: The repertory grid technique, a procedure based on a structured interview and a rating task, was administered to 68 early and late adolescents. Six measures of self-knowledge organization were computed from the data matrix obtained from each participant: differentiation; polarization; discrepancies between actual self (A), ideal self (I) and others (O); and the presence of inner conflicts. Regression-based moderation analyses were performed considering gender, adolescence phase and the interaction gender*adolescence phase as predictors. The association of the presence of conflict with gender and adolescence phase was tested with non-parametrical tests.

Results: Gender main effects suggested higher A-I ($b = 0.16$, boot 95% CI: 0.09, 0.57) and A-O ($b = 0.12$, boot 95% CI: 0.01, 0.26) self-discrepancies for boys, and gender*adolescence phase interaction effects were significant for polarization ($b = -22.51$, boot 95% CI: -41.92, -3.10), A-I ($b = -0.35$, boot 95% CI: -0.64, -0.01) and A-O ($b = -0.27$, boot 95% CI: -0.54, -0.05). Conditional effect probes revealed that polarization was lower at late adolescence only for boys, A-I and A-O self-discrepancies were higher at late adolescence only for girls. Moreover, more girls presented conflicts at late than early adolescence [χ^2 (1, $N = 47$) = 4.850, $p = .028$] and boys scored higher than girls on A-I and A-O self-discrepancies at early adolescence.

Conclusions: Gender differences were revealed in early adolescence: boys showed both lower self-esteem (higher A-I) and identification with others (higher A-O). Additionally, diverse patterns of developmental differences were suggested, such that, at late adolescence, polarization was lower only for boys, and self-esteem (higher A-I) and identification (higher A-O) were lower and conflicts more likely only for girls. Mixed results for differentiation and I-O self-differentiation recommend further research.

What rejection looks like: the proximity contacts of rejected children in a therapeutic setting – a preliminary study

Augusto Parras^{1,2}, Brenda M.S. da Silva^{1,2,3}, Hugo Folgado^{1,2}, Guida Veiga^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, PT;

²Comprehensive Health Research Centre, Universidade de Évora, Évora, PT; ³Developmental and Educational Psychology, Leiden University, Leiden, NL

Keywords: Radio-frequency; Sensor data; Sociometrics; Inclusion; Interactions; Psychomotor intervention

Objective: Peer rejection during childhood is related to many negative outcomes in adolescence and adulthood. During group psychomotor intervention children are able to cooperate with their peers, observe their skills, and share playful and ludic experiences together with the therapist as a mediator of their interactions. However, even in this protected setting rejected children might still struggle to fit in with their peer group. This pilot study has two aims: examine whether proximity contacts of rejected children differ from their peers at the baseline of a group psychomotor intervention; examine if proximity contacts of rejected children change throughout the intervention. Considering the importance of proximity contacts for the social emotional development of children, gaining knowledge on this matter is important so psychomotor therapist can adjust their practice in order to promote more social participation for children who are rejected within the peer group.

Methods: 17 children (4 rejected and 13 non-rejected) from grade 2 who attended schools in Évora participated in this study. The data used in this study derived from 14 psychomotor intervention sessions (2x week with 40min duration). During the session each child wore a RFID sensor that recorded their proximity contacts with other peers during the whole session. A proximity contact was recorded when children were in proximity (< 1.5 meters) and facing each other within a 65° angle. Information regarding partners of interaction, duration of social contacts, and time in solitude were obtained. Peer status was obtained through a peer nomination procedure. All scores were standardized per class.

Results: The Mann Whitney U test showed no differences regarding proximity contacts of rejected children and their peers were found at the baseline of a group psychomotor intervention. The Mann Whitney U test showed that regarding the means throughout the whole intervention rejected children spent less time in groups ($U = 4$, $p = .05$), and more time alone in total ($U = 3.00$, $p = .02$). Additionally their times in solitude lasted longer ($U = 7$, $p = .03$). However, a Wilcoxon matched paired test showed no differences between the social contacts of rejected children at baseline and after 14 sessions, variables suggesting that the intervention did not influence proximity contacts.

Conclusions: While the proximity contacts of rejected children at the beginning of the intervention program did not differ from their peers, their overall results throughout the sessions show that rejected children struggle with being included in the peer group even in the context of group psychomotor interventions.

Although group psychomotor intervention allows children to cooperate and develop together, our preliminary findings suggest that a more directed approach to group cohesion may be necessary to promote social inclusion of rejected children within the peer group. Future with larger sample sizes are necessary to provide more concrete and reliable conclusions about proximity contacts of rejected children in psychomotor interventions.

Relations between Motor Competence and Spatial Exploration of the Playground in first cycle children

Augusto Parras Albuquerque^{1,2}, Hugo Folgado^{2,3}, Guida Veiga^{2,3}

¹Instituto de Investigação e Formação Avançada – Universidade de Évora, Évora, PT; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, PT; ³Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, PT

Keywords: Motor Competence; Motor Coordination Test for Children (KTK); Playground; Positional Data; Spatial Exploration

Objective: The research aimed to examine the relationship between children's motor competence and spatial exploration, measured through individual spatial-temporal displacements during recess, in first grade elementary school students.

Methods: Motor competence was assessed through the Motor Coordination Test for Children (Körperkoordinationstest Für Kinder - KTK) and the results were crossed with the indicators of spatial exploration, calculated through the distances of each student to their midpoint at recess, identified from positional data collected by individual GPS devices placed of the playground. The study had 52 participants (23 girls and 29 boys) with a mean age of 6.25 years ($SD \pm 0.44$ years), from three first grade classes, located in the same school group in Évora-Portugal. To calculate the relationships between the variables, the non-parametric correlation tests of Kendall's Tau_b and Spearman's R ρ were used.

Results: The results showed a negative correlation ($\rho = -0.287$; $\tau = -0.196$; $p < 0.05$), between General Motor Coordination Coefficient (GMC – KTK) ($M: 363.24 \pm 62.22$) and the spatial exploration, captured by the average distance from to each children midpoint (11.47 ± 3.81 m).

Conclusions: The results suggest a negative relationship between motor competence and spatial exploration during recess. The better children perform on motor tasks, the less they explore the space around. This finding, combined with other studies, might suggest that children with better motor competence might be more stable on their engagement in physical activities during recess.

Effect of a motor-cognitive intervention on depression in middle-aged and older patients with type 2 diabetes mellitus

Nilton Leite^{1,2}, José Marmeira^{1,2}, Soraia Ferreira^{1,2}, Armando Raimundo^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano – Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal

Keywords: Mental health, Depression, Dual-task exercise, Type 2 Diabetes Mellitus

Objective: The main purpose of this study was to investigate the effect of a short-term motor-cognitive intervention on depression in patients with type 2 diabetes mellitus.

Methods: A one-group design with repeated measures was used. Twenty-six middle-aged and older patients with type 2 diabetes mellitus (68.6 ± 6.2 years; diabetes duration of 14.8 ± 8.4 years) acted as their own controls. Patients underwent baseline testing twice, with a 4-week interval between tests (pretest 1 and pretest 2) before participating in an exercise programme for a total of 8-weeks (3 times per week; average duration of 55-60 minutes per session). They were retested immediately after the end of the training programme. Exercise intensity, enjoyment level of exercise, and adverse events were systematically recorded at the end of each session. Depression symptoms was measured with the Geriatric Depression Scale (GDS-15), a self-administered questionnaire developed and validated to measure depressive symptoms in the older population.

Results: Participants reported a mean subjective exercise intensity and enjoyment level of 10.7 ± 1.4 points (Borg scale ranging from 6 to 20 points) and 4.0 ± 0.8 (out of 5 possible points), respectively. No adverse events were reported. After the training programme, statistical tests showed no significant improvement in depression symptoms over time ($p > 0.017$, Bonferroni adjustment to compensate for multiple comparisons).

Conclusions: The multimodal intervention combining physical and cognitive stimulation was found to be safe and enjoyable. Regarding depression scores, the intervention did not lead to significant changes in middle-aged and older patients with type 2 diabetes mellitus.

A method to quantify postural control during attention task in preschoolers

Ana Rita Matias^{1,2}, Filipe Melo^{1,3}, Paulo Delgado², Orlando Fernandes^{1,2}

¹Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano – Universidade de Évora, Évora, Portugal; ²Comprehensive Health Research Centre, Universidade de Évora, Évora, Portugal. ³Faculty of Human Kinetics - Lisbon University

Keywords: Children; Postural assessment; CAS

Objective: The relation between motor tasks (such as handwriting) and attention are closely related. It is common that educators mention postures readjustment in children with attention difficulties, which is not surprising considering that maintaining posture requires attentional resources. The attention task is from Cognitive Assessment System (CAS) is structured into four areas: planning, attention, and successive and simultaneous processing. Therefore, it is important to understand postural control behavior during an attention task to better prepare child's environment during motor tasks. This study aims to relate the traditional variables of postural control with the attention level of children assessed by the CAS.

Methods: Sixty-seven children (5-6 years) performed an attention task (CAS) while standing on a force plate to measure the center of pressure (CoP) excursions. For the assessment of postural control, the children were standing on the force plate (Plux -Portugal). The displacement of the CoP was sampled at 100 Hz.

Results: The results of this study show that there is a significant negative association between all traditional parameters and the outcome of the attention task (TOTEX; -0.399; < .001; TOTEX_AP; -0.364; 0.002; TOTEX_ML; -0.424; < .001; A_AP; -0.376; 0.002; A_ML; -0.394; < .001; AREA_CC; -0.448; < .001; AREA_CE; -0.343; 0.004).

Conclusions: We can conclude that the better the outcome in the task, the lower the values in the postural control parameters. This procedure will allow a deeper analysis of attention during motor tasks. The inclusion of other nonlinear measures will allow us to assess the complexity and regularity of the behavior of sway during these motor tasks.

The main concerns of people with Chronic Kidney Disease in disease management: a systematic review

Ana Rita Pedro^{1,2}, Beatriz Raposo^{2,3}, Fernando Genovez de Avelar^{2,3}, Ana Soraia Cunha^{2,3}

¹NOVA National School of Public Health, Public Health Research Center, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Lisbon, Portugal; ²NOVA National School of Public Health, NOVA University Lisbon, Lisbon, Portugal; ³NOVA National School of Public Health, Public Health Research Center, CISP, NOVA University Lisbon, Lisbon, Portugal

Keywords: Chronic Kidney Disease; Concerns; Patient-centered care; Patient experience; Systematic review

Objective: This study aimed to identify in a comprehensive way concerns of people with Chronic Kidney Disease (CKD) at different stages of this disease.

Methods: A systematic review was conducted according to PRISMA Statement Guidelines. Were searched qualitative, quantitative and mixed-methods studies in 4 databases (Scopus, Web of Science, PubMed and SCIELO) consulted in January '22 using established criteria, to answer the following research question: What are the concerns of people with CKD, in managing their disease? The protocol of this review was registered and published in PROSPERO (CRD42022308154). A methodological quality assessment was conducted using MMAT (Mixed Methods Appraisal Tool) version 2018. The results were integrated into a thematic synthesis.

Results: A total of 252 articles were identified, of these 10 met the methodological quality standards and inclusion criteria. The included studies were published between 2003 and 2020, originated from different countries and have a qualitative approach. The thematic synthesis identified 4 dimensions or general themes of concerns of people with CKD, namely: (i) concerns about treatment; (ii) concerns related to general health; (iii) concerns related with personal life and (iv) concerns related to self-care. A total of 37 items of concerns were identified from these dimensions, being the most identified: "fear of dying", "fear of not receiving the kidney transplant", "becoming ill or have complications before transplantation" and "fear of starting dialysis" and "change in lifestyle".

Conclusions: The most prevalent themes identified in this systematic review are directly related to the end-stage renal disease (ESRD), which reinforces the concentration of evidence at this stage. The available evidence on the concerns of people with CKD is still limited, which is an important barrier to be overcome for a better understanding of CKD and hence the effectiveness of health-focused actions, particularly in the early stages of the disease.